

ARTICLE OPEN

Investigating the association between obesity and asthma in 6- to 8-year-old Saudi children: a matched case–control study

Mahmoud Nahhas¹, Raj Bhopal², Chantelle Anandan¹, Rob Elton³ and Aziz Sheikh^{1,4}

BACKGROUND: Previous studies have demonstrated an association between obesity and asthma, but there remains considerable uncertainty about whether this reflects an underlying causal relationship.

AIMS: To investigate the association between obesity and asthma in pre-pubertal children and to investigate the roles of airway obstruction and atopy as possible causal mechanisms.

METHODS: We conducted an age- and sex-matched case–control study of 1,264 6- to 8-year-old schoolchildren with and without asthma recruited from 37 randomly selected schools in Madinah, Saudi Arabia. The body mass index (BMI), waist circumference and skin fold thickness of the 632 children with asthma were compared with those of the 632 control children without asthma. Associations between obesity and asthma, adjusted for other potential risk factors, were assessed separately in boys and girls using conditional logistic regression analysis. The possible mediating roles of atopy and airway obstruction were studied by investigating the impact of incorporating data on sensitisation to common aeroallergens and measurements of lung function.

RESULTS: BMI was associated with asthma in boys (odds ratio (OR) = 1.14, 95% confidence interval (CI), 1.08–1.20; adjusted OR = 1.11, 95% CI, 1.03–1.19) and girls (OR = 1.37, 95% CI, 1.26–1.50; adjusted OR = 1.38, 95% CI, 1.23–1.56). Adjusting for forced expiratory volume in 1 s had a negligible impact on these associations, but these were attenuated following adjustment for allergic sensitisation, particularly in girls (girls: OR = 1.25; 95% CI, 0.96–1.60; boys: OR = 1.09, 95% CI, 0.99–1.19).

CONCLUSIONS: BMI is associated with asthma in pre-pubertal Saudi boys and girls; this effect does not appear to be mediated through respiratory obstruction, but in girls this may at least partially be mediated through increased risk of allergic sensitisation.

npj Primary Care Respiratory Medicine (2014) **24**, Article number: 14004; doi:10.1038/npjpcrm.2014.4; published online 5 June 2014

INTRODUCTION

The substantial parallel increases in the prevalence of obesity^{1–5} and asthma^{6–8} observed over recent decades have led to the suggestion that obesity may be causally implicated in the risk of developing asthma.^{9–11} This line of enquiry began with the study of female nurses enrolled in the Nurses' Health Study cohort, which found a strong, dose-dependent relationship between an increase in body mass index (BMI) and adult-onset asthma.¹² Investigations into the association between obesity and asthma have subsequently been extended to both male and female adults,¹³ adolescents^{14–16} and young children,¹⁷ other populations of European origin^{9,18–20} and some other ethnic groups.¹⁷ Overall, this body of work has confirmed the original observation—namely that obesity is associated with an increased risk of developing asthma,¹⁵ particularly in post-pubertal females.²¹ What is, however, far less clear is whether this reflects a causal relationship between the two and, if so, what the underlying mechanisms might be.

Judgements on causal inference can be greatly aided by studying biological mechanisms²² and it is therefore welcome that the epidemiological evidence has triggered a substantial body of mechanistic work in both animal and human populations.^{23,24} Broadly speaking, this work has converged on studying the role of common genes,^{25,26} fetal programming,²⁷ sex hormones,^{11,27,28} respiratory obstruction,²⁹ bronchial hyper-reactivity²¹ and changes

in immunological and inflammatory responses.^{11,26,27} This body of work has, however, thus far failed to generate any clear insights into possible biological pathways, possibly reflecting the fact that different mechanisms may be implicated in different population subgroups. It has therefore been suggested that age- and gender-specific investigations be undertaken in an attempt to progress understanding on possible causal pathways.²¹

We sought to contribute to this important area of enquiry by undertaking the first study of obesity and asthma in a Middle Eastern population. We investigated the association between obesity and asthma in pre-pubertal school-aged children, undertaking separate analyses in boys and girls, with a view to understanding the possible aetiological roles of physiological (i.e., respiratory obstruction) and immunological (i.e., allergic sensitisation) mechanisms.

MATERIALS AND METHODS

Ethical considerations

Permission was obtained from the education authorities in Madinah (General Directors) and School Health Departments in the Ministry of Education, Riyadh, Saudi Arabia, and from head-teachers of participating schools. Parents gave their signed, informed consent to their child's participation. All data had identifiers removed to minimise the risk of inadvertently breaching confidentiality.

¹Allergy and Respiratory Research Group, Centre for Population Health Sciences, Medical School, The University of Edinburgh, Edinburgh, UK; ²Bruce and John Usher Professor of Public Health, Centre for Population Health Sciences, Medical School, The University of Edinburgh, Edinburgh, UK; ³Centre for Population Health Sciences, Medical School, The University of Edinburgh, Edinburgh, UK and ⁴Division of General Internal Medicine and Primary Care, Brigham and Women's Hospital/Harvard Medical School, Boston, MA, USA. Correspondence: A Sheikh (aziz.sheikh@ed.ac.uk)

Received 29 September 2013; revised 28 February 2014; accepted 1 March 2014

Design

We conducted an age- and sex-matched case-control study of 1,264 schoolchildren with and without asthma. Children aged 6–8 years were recruited from 37 randomly selected schools in Madinah, Saudi Arabia. These children were identified from our linked cross-sectional study of over 5,000 children in whom we found an asthma prevalence of 23.6% (95% confidence interval (CI), 21.3, 26.0); this group of children provided a sampling frame for this investigation.³⁰ For this phase of work, we compared BMI, triceps skin fold thickness (TSFT) and waist circumferences (WCs) in children with a history of symptoms indicative of asthma (cases) with individually age- and sex-matched children without such a history (controls), factoring respiratory function and allergic sensitisation into the analyses. Pilot work confirmed the feasibility of undertaking this research.

Setting

Madinah is a city in the west of Saudi Arabia, with an area of 589 km², which lies approximately 190 km from the Red Sea coast. It has a population of over 1.5 million people, approximately 5% of the total of 21.4 million residents in Saudi Arabia.³¹

Recruitment of schools and students

Lists of boys' and girls' primary schools were obtained from the General Directorate of Education of the Madinah region. Schools were then stratified according to the five geographical areas of Madinah and students' gender and a random sample of 37 schools (i.e., 8 girls' and 29 boys' schools) were approached. We invited the schools to participate with a letter of explanation to the head-teachers outlining the purpose of the study and the procedures that were to be employed.

Eligible children were those aged 6–8 years who had been resident in Madinah for at least 1 year prior to the start of the study. All eligible subjects were invited to participate via a letter and a consent form sent by the school to their parents.

Identification of cases and controls

The parents of all recruited children were asked to complete a questionnaire incorporating validated questions from the International Study of Asthma and Allergies in Childhood (ISAAC)³² that had previously been adapted for use in Arab populations.³³ Cases were defined as children who were identified by parents as those who had 'ever had wheeze',^{30,34–36} controls were defined as those children, matched for age and sex, who had according to their parents 'never had wheeze' as reported using the same instrument.

Measures of obesity

Our primary measure of obesity was BMI,³⁷ but, in addition, we measured weight, TSFT and WC. Children were asked to wear normal light clothing during measurements and to take off their shoes. Measurements of height to enable calculation of BMI (which was calculated as weight (kg) divided by the square of height in metres (kg/m²)), weight, TSFT and WC were recorded in the schools by trained members of the health-care team who were unaware of whether children had asthma or not. Height was measured with a portable stadiometer (SECA Leicester height measure, SECA, Birmingham, UK) using the method described by Tanner *et al.*³⁸ Measurements were made to the nearest 0.1 cm.

Children were weighed and measured separately from their classmates. Weight was measured using SECA Mechanical Column Scales, with measurements being recorded to the nearest 0.1 kg.

We used BMI percentiles for girls and boys per age as a measure of standardised weight. As there are no reference data available for Saudi children, we used the Center for Disease Control and Prevention (CDC) BMI-for-age growth charts for girls and boys.³⁹ According to these charts, underweight was defined as a BMI less than the 5th centile, healthy weight was defined as a BMI between the 5th and the 85th centile, overweight was defined as a BMI between the 85th and 94th centiles and obese was defined as a BMI equal to or greater than the 95th centile.

TSFT was measured three times to the nearest millimetre with a Holtain Skinfold Caliper (Holtain, Crymych, UK) on the right arm. The triceps skin fold locus was defined as being halfway between the acromion and olecranon on the back of the arm measured with the elbow bent.^{40,41} TSFT was used mainly to determine relative obesity and the percentage of body fat.^{40,41}

WC was measured to the nearest centimetre with a Lufkin flexible steel tape measure (Apex Tool Group, Sparks, MD, USA), with children in the

standing position after gentle expiration. The following anatomical landmarks were used: laterally, midway between the lowest portion of the rib cage and iliac crest; and anteriorly, midway between the xiphoid process of the sternum and the umbilicus.⁴²

Potential confounders

The parental questionnaire provided information on the following potential confounders:

- Parental age
- Parental education
- Parental smoking status
- Number of siblings
- Place of birth
- Breastfeeding
- Medications administered (i.e., antibiotics and paracetamol)
- Exposure to pets and farm animals
- Food and other lifestyle factors

Measurement of explanatory variables

We measured lung function and undertook skin prick tests to common aeroallergens to investigate whether the presence of respiratory obstruction or allergic sensitisation could help explain any possible associations uncovered.

Lung function tests. Lung function testing was performed in schools by trained members of the research team. All spirometric measurements were taken in a sitting position using a Vitalograph Pneumotrac spirometer (Vitalograph, Buckingham, UK). England The best of at least three technically acceptable values for forced expiratory volume in 1 s (FEV₁), forced vital capacity (FVC), (FEV₁%) FEV₁/FVC ratio and peak expiratory flow were selected.^{43–45}

Skin prick tests. Skin prick tests were undertaken using the volar aspect of the forearm, the skin having first been marked with a pen. Thereafter, one drop of each solution to be tested was put on the skin beside the respective mark. A panel of eight allergens (i.e., cat, *Dermatophagoides farinae*, *Dermatophagoides pteronyssinus*, grass mix, Rye, tree mix (early blossoming), tree mix (mid-blossoming) and weed mix) were used. Histamine 10 mg/ml and 50% glycerine were used as positive and negative controls, respectively. The size of the weal was recorded as the mean of the greatest diameter and the diameter perpendicular to its midpoint. Recordings were taken after 15 min and a mean diameter of 3 mm or greater than the negative control was regarded as positive.⁴⁶

Statistical methods

All statistical analyses were carried out using Statistical Package for Social Sciences (SPSS, Chicago, IL, USA), version 16.0, for Windows and Confidence Interval Analysis (CIA) software (University of Southampton, Southampton, UK).⁴⁷ All analyses were undertaken separately for boys and girls because the literature suggested that we might find differing degrees of association between obesity and asthma.^{21,28} Univariate comparisons between the matched cases and controls were carried out by paired *t*-test variables and Wilson's method for binary variables. The differences in measures of obesity between the matched pairs of children were tested for variation between schools, separately for each sex, by one-way analysis of variance. There were no significant differences between schools in any of the four measures, suggesting no important cluster effects, and all further analyses were therefore carried out without any adjustment for school effects. Multiple conditional logistic regression was used to investigate the relationship between asthma and explanatory variables, adjusted for confounders. In the regression models, only those variables that differed significantly between cases and controls were included as covariates.

Sample size calculations

Our preliminary work³⁰ suggested that the prevalence of asthma was around 15% and therefore a sample of around 5,000 children needed to be recruited into the cross-sectional study in order to recruit sufficient numbers of cases (≥ 600) to give 80% power to detect a mean difference of around 0.4 in BMI between cases and the same number of controls based on a s.d. of three units for BMI.

RESULTS

Recruitment of schools

All 37 schools that were approached agreed to participate.

Characteristics of cases and controls

Data were collected on 388 matched pairs of boys and 244 matched pairs of girls. Table 1 summarises the main characteristics of cases and controls among boys and girls. The demographic and lifestyle variables that differed between cases and controls in males were birth weight, smoking habits of the father, breastfeeding, taking of paracetamol and antibiotics during the first year of life, taking of paracetamol in the previous 12 months and the frequency of trucks passing near the home. In girls, demographic

and lifestyle variables that differed between cases and controls were smoking habits of the father, breastfeeding, exposure to farm animals during the first year of life, taking paracetamol during the previous 12 months, exposure to a cat during the previous 12 months, the type of air conditioning system at home and meat and nut consumption.

Univariate analysis

Table 2 shows the unadjusted effect of each of the obesity measures on the risk of having asthma in boys and girls. All four obesity measures (BMI, weight, TSFT and WC) were found to be positively associated with asthma in both boys and girls. The odds ratios in girls were consistently higher than those in boys.

Table 1. Descriptive comparisons between cases and controls in male and female children: figures shown for cases and controls are mean (s.d.) or number (%)

Variables	Males			Females		
	Cases (n = 388)	Controls (n = 388)	Differences for the mean or for rates between cases and controls (95% CI)	Cases (n = 244)	Controls (n = 244)	Differences for the mean or for rates between cases and controls (95% CI)
Measures of obesity						
Primary measures mean (s.d.)						
BMI (kg/m ²)	17.0 (3.8)	15.8 (2.5)	1.2 (+0.7, +1.7) ^a	19.0 (4.4)	15.9 (2.1)	3.1 (+2.5, +3.7) ^a
Secondary measures mean (s.d.)						
TSFT (mm)	10.5 (4.2)	9.2 (3.3)	1.3 (+0.8, +1.8) ^a	13.8 (5.3)	10.5 (3.2)	3.3 (+2.6, +4.1) ^a
WC (cm)	56.6 (8.8)	53.6 (6.4)	3 (+1.9, +4.0) ^a	61.0 (10.1)	53.9 (5.8)	7.1 (+5.6, +8.4) ^a
Weight (kg)	26.1 (8.0)	23.9 (5.7)	2.2 (+1.2, +3.2) ^a	30.4 (9.6)	24.1 (5.4)	6.3 (+4.92, +7.69) ^a
Explanatory variables						
Lung function measures						
FVC (Lt) mean (s.d.)	1.35 (0.36)	1.46 (0.37)	-0.11 (-0.16, -0.06) ^a	1.17 (0.31)	1.22 (0.30)	-0.05 (-0.10, +0.00)
FEV ₁ (Lt) mean (s.d.)	1.15 (0.30)	1.31 (0.34)	-0.16 (-0.21, -0.12) ^a	0.96 (0.27)	1.08 (0.28)	-0.12 (-0.17, -0.08) ^a
PEFR (Lt) mean (s.d.)	2.40 (0.64)	2.72 (0.56)	-0.32 (-0.41, -0.24) ^a	1.77 (0.76)	2.36 (0.57)	-0.59 (-0.70, -0.47) ^a
Sensitisation to one or more allergens (≥3 mm), n (%)						
Cat	216 (56)	55 (14)	42 (+36.0, +48.0) ^a	166 (68)	35 (14)	54 (+46.7, +61.3) ^a
Dermatophagoides farinae	99 (25)	21 (5)	20 (+15.2, +24.8) ^a	79 (32)	6 (2)	30 (+23.9, +36.1) ^a
Dermatophagoides pteronyssinus	99 (25)	11 (3)	22 (+17.4, +26.6) ^a	91 (37)	16 (6)	31 (+24.3, +37.8) ^a
Grass mix	66 (17)	14 (4)	13 (+8.8, +17.2) ^a	51 (21)	5 (2)	19 (+13.6, +24.4) ^a
Rye	55 (14)	15 (4)	10 (+6.0, +14.0) ^a	83 (34)	2 (1)	33 (+26.9, +39.1) ^a
Tree mix (early blossoming)	70 (18)	13 (3)	15 (+10.8, +19.2) ^a	89 (36)	4 (2)	32 (+27.7, +40.3) ^a
Tree mix (mid-blossoming)	80 (21)	18 (5)	16 (+11.4, +20.6) ^a	81 (33)	3 (1)	30 (+26.0, +38.0) ^a
Weed mix	62 (16)	10 (3)	13 (+9.0, +17.0) ^a	66 (27)	7 (2)	20 (+19.2, +30.8) ^a
Sensitisation to 1 ≥ allergen	325 (84)	107 (27)	57 (+51.3, +62.7) ^a	234 (96)	54 (22)	74 (+68.2, +79.8) ^a
Potential confounders						
Father's age (years) mean (s.d.)	43.3 (8.4)	42.1 (7.8)	1.12 (-0.04, +2.28)	44.1 (9.0)	44.8 (8.9)	0.1 (-5.0, +5.2)
Mother's age (years) mean (s.d.)	35 (6.1)	34.6 (6.2)	0.36 (-0.51, +1.23)	36.2 (6.0)	36.1 (6.1)	-0.1 (-4.3, +4.1)
Birth weight (kg) mean (s.d.)	2.90 (0.66)	3.00 (0.58)	-0.10 (-0.20, -0.01) ^a	2.94 (0.67)	2.83 (0.62)	0.11 (-0.02, +0.24)
Father's education (highest qualification), n (%)						
None	14 (3.7)	21 (5.6)	-1.9 (-4.9, +1.1)	22 (9.4)	14 (5.9)	3.5 (-1.2, +8.2)
General education	192 (50.8)	198 (53.2)	-2.4 (-1.0, +4.0)	132 (56.2)	156 (65.3)	-9.1 (-8.9, +8.7)
Higher education	172 (45.5)	153 (41.1)	4.4 (-3.0, +11.0)	81 (34.5)	69 (28.9)	5.6 (-2.6, +13.6)
Mother's education (highest qualification), n (%)						
None	14 (3.7)	32 (8.3)	-4.6 (-7.9, -1.3) ^a	26 (10.9)	29 (12.0)	-1.1 (-6.7, +4.5)
General education	208 (54.5)	205 (53.4)	1.1 (-5.9, +8.1)	141 (59.0)	144 (59.8)	-0.8 (-9.5, +7.9)
Higher education	160 (41.9)	147 (38.3)	3.6 (-3.3, +10.5)	72 (30.1)	68 (28.2)	1.9 (-6.2, +10)
Father smoker, n (%)						
Yes	129 (33.3)	96 (24.8)	8.5 (+2.1, +14.9) ^a	56 (23.1)	77 (31.8)	-8.7 (-16.6, -0.8) ^a
No	258 (66.7)	291 (75.2)	-8.5 (-14.9, -2.1) ^a	186 (76.9)	168 (68.2)	8.7 (+0.8, +16.6) ^a
Mother smoker, n (%)						
Yes	5 (1.3)	1 (0.3)	1.0 (-0.25, +2.25)	4 (1.7)	2 (0.8)	0.9 (-1.1, +2.9)
No	382 (98.7)	385 (99.7)	-1.0 (-2.3, +0.3)	236 (98.3)	241 (99.2)	-0.9 (-2.9, +1.1)
Birth order, n (%)						
First child	206 (53.4)	206 (53.1)	0.3 (-6.7, +7.3)	43 (17.8)	47 (19.5)	-1.7 (-8.6, +5.2)
Second child	125 (32.4)	123 (31.7)	0.7 (-5.9, +7.3)	189 (78.1)	180 (74.7)	3.4 (-4.1, +10.9)
Third or greater	55 (14.2)	56 (15.2)	-1.0 (-6.0, +4.0)	10 (4.1)	14 (5.8)	-1.7 (-5.5, +2.1)
Was the child born in Madinah?, n (%)						
Yes	332 (85.8)	328 (85.4)	0.4 (-4.5, +5.3)	202 (84.5)	200 (82.6)	1.9 (-4.7, +8.5)
No	55 (14.2)	56 (14.6)	-0.4 (-5.3, +4.5)	37 (15.5)	42 (17.4)	-1.9 (-8.5, +4.7)
Did the mother breastfeed?, n (%)						
Yes	290 (75.9)	330 (85.5)	-9.6 (-15.1, -4.1) ^a	208 (86.7)	193 (80.1)	6.6 (0.0, +13.0)
No	92 (24.1)	56 (14.5)	8.5 (+2.1, +14.9) ^a	32 (13.3)	48 (19.9)	-8.7 (-16.6, -0.8) ^a
Medication given in the 1st year of child's life, n (%)						
Paracetamol	367 (95.6)	329 (88.4)	7.2 (+3.4, +11.0) ^a	213 (89.9)	209 (87.4)	2.5 (-3.1, +8.1)
Antibiotics	311 (82.3)	213 (57.9)	24.4 (+18.2, +30.6) ^a	164 (69.5)	148 (61.7)	7.8 (-0.6, +16.2)

Table 1. (Continued)

Variables	Males			Females		
	Cases (n = 388)	Controls (n = 388)	Differences for the mean or for rates between cases and controls (95% CI)	Cases (n = 244)	Controls (n = 244)	Differences for the mean or for rates between cases and controls (95% CI)
Paracetamol administered in the last 12 months, n (%)						
Once a week at least	87 (22.9)	70 (18.9)	4.0 (−1.7, +9.7)	54 (22.9)	39 (16.9)	6.0 (−1.1, +13.1)
Once a month at least	220 (57.9)	195 (52.7)	5.2 (−1.8, +12.2)	135 (57.2)	125 (54.1)	3.1 (−5.7, +11.9)
Once a year at least	73 (19.2)	105 (28.4)	−9.2 (−15.2, −3.2) ^a	47 (19.9)	67 (29.0)	−9.1 (−16.7, −1.5) ^a
Exposure to animals, n (%)						
Farm animal when mother was pregnant	11 (2.9)	11 (2.9)	0.0 (−2.4, +2.4)	14 (6.1)	8 (3.4)	2.7 (−1.1, +6.5)
Cat at home in the 1st year of child's life	15 (3.9)	15 (3.9)	0.0 (−2.7, +2.7)	19 (7.9)	6 (2.5)	5.4 (−1.5, +9.3) ^a
Cat at home in the last 12 months	29 (7.5)	25 (6.5)	1.0 (−2.6, +4.6)	18 (7.5)	12 (5.0)	2.5 (−1.2, +8.8) ^a
Farm animal in the 1st year of child life	24 (6.2)	16 (4.2)	2.0 (−1.1, +5.1)	16 (6.8)	5 (2.1)	4.7 (−1.1, +8.3) ^a
Taking exercise per week, n (%)						
Never	241 (64.3)	254 (68.8)	−4.5 (−11.1, +2.1)	201 (85.5)	200 (86.2)	−0.7 (−6.9, +5.5)
Once or twice a week	101 (26.9)	86 (23.3)	3.6 (−2.5, +9.7)	31 (13.2)	25 (10.8)	2.4 (−3.4, +8.2)
Three or more a week	33 (8.8)	29 (7.9)	0.9 (−3.0, +4.8)	3 (1.3)	7 (3.0)	−1.7 (−4.3, +0.9)
Watching TV, n (%)						
< 3 h per day	228 (60.0)	241 (64.6)	−4.6 (−11.4, +2.2)	137 (56.8)	128 (54.9)	1.9 (−6.9, +10.7)
> 3 h per day	152 (40.0)	132 (35.4)	4.6 (−2.2, +11.4)	104 (43.2)	105 (45.1)	−1.9 (−10.7, +6.9)
What is the fuel normally used in cooking in your household?, n (%)						
Electricity only	20 (5.2)	20 (5.2)	0 (−3.1, +3.1)	9 (3.8)	15 (6.3)	−2.5 (−6.4, +1.4)
Gas only	351 (90.9)	349 (90.4)	0.5 (−3.6, +4.6)	223 (92.9)	220 (92.4)	0.5 (−4.1, +5.1)
Both electricity & gas	15 (3.9)	17 (4.4)	−0.5 (−3.3, +2.3)	8 (3.3)	3 (1.3)	2.0 (−0.7, +4.7)
Air conditioning type, n (%)						
Electric fan only	3 (0.8)	1 (0.3)	0.5 (−0.54, +1.54)	1 (0.4)	2 (0.8)	−0.4 (−1.77, +0.97)
Water system only	4 (1.0)	5 (1.3)	−0.3 (−1.8, +1.2)	4 (1.7)	10 (4.1)	−2.4 (−5.4, +0.6)
Freon system only	365 (94.3)	372 (96.6)	−2.3 (−5.2, +0.6)	194 (82.2)	223 (92.5)	−10.3 (−16.1, −4.5) ^a
Both Freon & water systems	15 (3.9)	7 (1.8)	2.1 (−0.2, +4.4)	37 (15.7)	6 (2.5)	13.2 (−8.2, +18.2) ^a
How often does a truck pass through the street adjacent to your home?, n (%)						
Rarely	237 (61.6)	268 (70.9)	−9.3 (−15.9, −2.7) ^a	168 (70.3)	169 (70.7)	−0.4 (−8.5, +7.7)
Frequently	148 (38.4)	110 (29.1)	9.3 (−2.7, +15.9) ^a	71 (29.7)	70 (29.3)	0.4 (−7.7, +8.5)
Diet—How many times a week does your child eat the following?, n (%)						
Meats						
Never	20 (5.2)	16 (4.2)	1.0 (−4.0, +2.0)	9 (3.7)	26 (10.9)	−7.2 (−11.8, −2.6) ^a
Once or twice a week	110 (28.6)	122 (31.8)	−3.2 (−9.7, +3.3)	83 (34.2)	81 (33.9)	0.3 (−8.1, +8.7)
Three or more a week	254 (66.1)	246 (64.1)	2.0 (−4.7, +8.7)	151 (62.1)	132 (55.2)	6.9 (−1.8, +15.6)
Fruits						
Never	68 (17.7)	44 (11.5)	6.2 (−6.9, +19.3)	29 (12.0)	24 (10.0)	2.0 (−3.5, +7.5)
Once or twice a week	167 (43.4)	175 (45.9)	−2.5 (−9.5, +4.5)	114 (47.1)	115 (48.1)	−1.0 (−9.9, +7.9)
Three or more a week	150 (39.0)	162 (42.5)	−3.5 (−10.4, +3.4)	99 (40.9)	100 (41.8)	−0.9 (−9.6, +7.8)
Rice						
Never	9 (2.4)	8 (2.1)	0.3 (−1.8, +2.4)	6 (2.5)	10 (4.2)	−1.7 (−4.9, +1.5)
Once or twice a week	70 (18.4)	83 (21.7)	−3.3 (−8.9, +2.3)	45 (19.0)	47 (19.7)	−0.7 (−7.7, +6.3)
Three or more a week	301 (79.2)	291 (76.2)	3.0 (−2.9, +8.9)	186 (78.2)	182 (76.2)	2.0 (−5.4, +9.4)
Nuts						
Never	224 (59.4)	227 (60.7)	−1.3 (−8.2, +5.6)	151 (62.9)	116 (50.0)	12.9 (−4.2, +21.6) ^a
Once or twice a week	134 (35.5)	118 (31.6)	3.9 (−2.7, +10.5)	73 (30.4)	93 (40.1)	−9.7 (−18.1, −1.3) ^a
Three or more a week	19 (5.0)	29 (7.8)	−2.8 (−6.2, +0.6)	16 (6.7)	23 (9.9)	−3.2 (−8.1, +1.7)
Fast food						
Never	217 (56.7)	225 (59.2)	−2.5 (−9.4, +4.4)	160 (67.5)	164 (69.8)	−2.3 (−10.5, +5.9)
Once or twice a week	132 (34.5)	122 (32.1)	2.4 (−4.2, +9.0)	60 (25.3)	56 (23.8)	1.5 (−6.1, +9.1)
Three or more a week	34 (8.9)	33 (8.7)	0.2 (−3.8, +4.2)	17 (7.2)	15 (6.4)	0.8 (−3.7, +5.3)

Note: Missing data for individual variables are not shown.

Abbreviations: BMI, body mass index; CI, confidence interval; TSFT, triceps skin fold thickness; WC, waist circumference.

^aStatistically significant difference between cases and controls.

Multivariate analysis

Table 3 shows the effect of BMI on the risk of asthma in boys and girls, adjusted for relevant potential confounding sociodemographic and lifestyle variables. Only those variables showing univariate associations with asthma were included in each model. These analyses showed that BMI was still strongly associated with asthma after adjustment for these covariates in both boys and girls. The adjusted odds ratio was higher in girls than in boys.

Investigation of possible causal pathways

There was—in both sexes—strong evidence that children with asthma had greater respiratory obstruction and were more sensitised to aeroallergens in comparison with those with no history of asthma. Table 4 shows the impact of factoring measures of respiratory obstruction and this reveals a negligible impact on the associations between BMI and asthma in boys and girls.

Table 2. Unadjusted effect of obesity and adiposity measures (TSFT, WC and weight) on the odds of asthma in males and females

Measures of obesity and adiposity	Multiplicative increase in odds of having asthma (95% CI) per unit increase of the independent variable	
	Males	Females
Obesity measure BMI (kg/m ²)	1.14 (1.08, 1.20)	1.37 (1.26, 1.50)
Adiposity measures		
TSFT (mm)	1.10 (1.06, 1.15)	1.24 (1.16, 1.33)
WC (cm)	1.06 (1.04, 1.08)	1.14 (1.10, 1.18)
Weight (kg)	1.06 (1.03, 1.08)	1.16 (1.12, 1.20)

Abbreviations: BMI, body mass index; CI, confidence interval; TSFT, triceps skin fold thickness; WC, waist circumference.

Table 3. Adjusted ORs (95% CI) for the association between BMI on asthma in males and females

Independent (X) variables in the multiple logistic regression model ^a	ORs of developing asthma (95% CI) adjusted for other factors	
	Males	Females
BMI	1.11 (1.03, 1.19)	1.38 (1.23, 1.56)
Birth weight	0.71 (0.49, 1.02)	^b
Mother education (none compared with higher)	0.48 (0.16, 1.44)	^b
Mother education (general compared with higher)	0.82 (0.51, 1.32)	^b
Father smoking	1.17 (0.76, 1.80)	0.45 (0.22, 0.93)
Breastfed	0.74 (0.44, 1.26)	3.62 (1.42, 9.25)
Paracetamol given in the 1st year of child's life	1.66 (0.67, 4.09)	^b
Antibiotics given in the 1st year of child's life	2.45 (1.54, 3.91)	^b
Paracetamol given to the child in the last 12 months (once a week compared with yearly)	1.94 (1.00, 3.80)	7.09 (2.48, 20.8)
Paracetamol given to the child in the last 12 months (once a month compared with yearly)	1.90 (1.14, 3.14)	2.23 (1.05, 4.72)
Cat at home in the 1st year of child's life	^c	0.83 (0.10, 7.28)
Cat at home in the last 12 months	^c	2.25 (0.42, 12.1)
Farm animal in the 1st year of child's life	^c	2.56 (0.36, 18.4)
Air conditioning type used at home (electric fan compared with Freon system)	^c	3.58 (0.02, 52.9)
Air conditioning type used at home (water system compared with Freon system)	^c	0.51 (0.09, 2.96)
Air conditioning type used at home (both Freon and water system compared with Freon system)	^c	19.4 (3.56, 106)
Frequency of a truck passing through the street adjacent to the home (frequently compared with rarely)	1.39 (0.89, 2.78)	^b
Eating meat (once or twice a week compared with never)	^c	10.2 (2.03, 51.2)
Eating meat (three times a week or more compared with never)	^c	8.68 (1.83, 41.0)
Eating nuts (once or twice a week compared with never)	^c	0.55 (0.29, 1.04)
Eating nuts (three times a week or more compared with never)	^c	0.80 (0.27, 2.37)

Abbreviations: BMI, body mass index; CI, confidence interval; OR, odds ratio.

^aOnly those variables that remained in the model in the last step of the backward stepwise regression process are displayed in the table.

^bNot in the model for females.

^cNot in the model for males.

Table 4. Estimates (95% CIs) for the coefficient of BMI in conditional logistic regression analysis adjusted for selected lung function and sensitisation and also for other factors found to be associated with obesity in Table 1

Lung and sensitisation factors	Males	Females
None	1.11 (1.03, 1.19)	1.38 (1.23, 1.56)
FEV ₁	1.10 (1.02, 1.18)	1.37 (1.22, 1.54)
FVC	1.11 (1.03, 1.19)	1.40 (1.24, 1.57)
Sensitisation to more than one allergen	1.09 (0.99, 1.19)	1.25 (0.96, 1.60)
FEV ₁ +sensitisation to more than one allergen	1.09 (0.98, 1.19)	1.28 (0.96, 1.69)
FVC+sensitisation to more than one allergen	1.09 (0.99, 1.19)	1.25 (0.97, 1.61)

Abbreviations: BMI, body mass index; CI, confidence interval; FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity.

Table 4 also reveals the impact of factoring the presence of allergic sensitisation into the analysis and this in contrast shows a more marked reduction in the associations in girls, although a more modest impact in boys.

DISCUSSION

Main findings

This large case-control study has demonstrated consistent associations between measures of overall, central and peripheral obesity, and asthma in boys and girls. In keeping with previous work,^{17,20} we found a stronger association in girls than in boys,

but unlike other studies we observed this difference in the context of studying pre-pubertal children. Focusing on BMI, we then demonstrated that this association was still present after adjusting for a wide range of potential demographic and lifestyle-related confounding factors. In an attempt to investigate whether respiratory obstruction and/or allergic sensitisation may lie on the causal pathway between obesity and asthma,^{11,26} we adjusted separately and then combined for their effects and found that respiratory obstruction had only a very modest impact, but that the association was more attenuated when adjusting for sensitisation, particularly in girls. Adjusting simultaneously for both lung function and sensitisation did not further attenuate the relationship, suggesting that respiratory obstruction is unlikely to be an important mechanism but that obesity may at least partially exert its effects by increasing the risk of allergic sensitisation. Overall, this study therefore adds to the increasing body of evidence implicating obesity as a risk factor for the development of asthma showing that this relationship also holds true in a population of Middle Eastern origin and provides important pointers into how this relationship may or may not be mediated in this population.

Strengths and limitations of this study

Key strengths of this work include the fact that we conducted a large, adequately powered, investigation, our use of a validated and extensively used questionnaire to identify cases and controls, careful measurements of a range of measures of obesity¹⁷ by trained personnel who were blinded to case status the measurement of, and adjustments for, a range of potential confounders and the *a priori* decision to undertake separate analyses in boys and girls.^{21,28} In addition, we sought to extend previous work by investigating two important possible causal mechanisms.

The main limitations of this work are those inherent in case-control studies—namely the risk of residual confounding, recall bias and an inability to establish a temporal relationship between the onset of obesity and the subsequent development of asthma; reverse causality therefore remains a possibility. Moreover, the identification of cases was based on responses to the ISAAC questionnaire and this may have been affected by parental behaviours and diagnostic bias; although a validated culturally appropriate questionnaire was used, there remains therefore the possibility of misclassification errors. If present, this is likely to have been operating in a non-differential manner, which would have resulted in blunting the associations between obesity and asthma rather than creating spurious associations. Future work should seek to build on this by using more objective assessments to diagnose asthma, although given the numbers of subjects who need to be studied this may well prove financially and logistically challenging.

Interpretation of findings in relation to previously published research

Our study is consistent with an increasing number of epidemiological studies that have reported a positive association between obesity and the risk of developing childhood asthma.^{17,20} As with other investigators,¹⁷ we have found that this association was stronger in girls. Whether this sex difference is simply due to the higher levels of obesity observed in female cases (Table 1) or due to difference in body fat distribution—particularly centrally, where it is believed to exert the greatest inflammatory effect,^{48,49} as reflected by the larger WC and prevalence of sensitisation in females with asthma (Table 1)—and/or associated biochemical responses is unclear.^{48,49} The fact that sensitisation was a strong risk factor for asthma confirms the findings of a number of previous epidemiological investigations.⁴⁶

There is a limited body of evidence investigating the relationship between obesity and allergic sensitisation, and the available evidence has been noted to be inconsistent and inconclusive.²¹ A large and important recent Swedish birth cohort study found that increased BMI at age 7 was associated with an increased risk of allergic sensitisation to aeroallergens and the development of childhood asthma, but that asthma risk was not increased if the increased BMI was confined to the first 4 years of life.⁵⁰ It is thus likely that any possible relationship between obesity and allergic sensitisation is strongly influenced by age. Sex is also important as recent work has also demonstrated that levels of the adipocyte-derived hormone leptin are closely correlated with BMI in early life, and that leptin levels tend to be higher in girls than in boys,⁵¹ which may explain why a stronger and more consistent relationship is observed between BMI and asthma in girls.²¹ Future work should now aim to study carefully characterised populations and also progress mechanistic lines of enquiry through measurement of key biomarkers including the adipokines adiponectin and leptin, other obesity-related hormones such as insulin and neuropeptides, and the pro-inflammatory cytokines tumour necrosis factor, interleukin-6 and interleukin-1B;²⁷ it is also important to investigate whether obesity reduces immunological tolerance through downregulation of interleukin-10 secretion and study the combined impact of these changes on regulatory T lymphocytes (Tregs).⁵²

Implications for future research, policy and practice

The findings of the present study may be valuable for public health authorities as it provides additional evidence of the considerable adverse risks associated with obesity. Although causality cannot be proven by this or indeed any other epidemiological investigation, the findings from this work, when combined with an increasing number of studies worldwide,^{12,13,17,20} provide increasing evidence of a causal relationship between obesity and asthma. This then

adds weight to the argument to consider and investigate the role of interventions to prevent/tackle obesity early on in life and thereby possibly reduce the risk of developing asthma. Even if such causality has not been unequivocally proven, public health authorities should consider applying the 'precautionary principle'⁵³ and undertake initiatives for reducing obesity among children. Reduction of obesity in children will offer benefits in the general health status, as childhood obesity is a well-established risk factor for several other diseases.⁵⁴ Moreover, given the high prevalence of obesity and asthma, school health departments should consider focussing attention on establishing systems for monitoring obesity and asthma among schoolchildren.

CONCLUSIONS

In conclusion, our large matched case-control study provides the first evidence demonstrating an association between obesity and asthma in pre-pubertal male and female Saudi children. Our investigation of key causal pathways suggests that this relationship may indeed be causal, being at least partially mediated through the pro-inflammatory effects of central obesity, which increase the risk of allergic sensitisation to aeroallergens. This work has also demonstrated that mechanical obstruction is unlikely to be an important aetiological mechanism in overweight, pre-pubertal children.

ACKNOWLEDGEMENTS

We thank the General Directorate of Education in Madinah region, the General Directorate of School Health in the Ministry of Education in Riyadh (both boys and girls sections), the research team, and all staff in the participating schools. We also thank the parents and children who participated in this study.

CONTRIBUTIONS

AS, MN, RB and CA were involved in conceiving and designing this study. MN undertook the field work under the supervision of AS and analysed the data under the supervision of RE. MN and AS led the drafting of the manuscript and RB, CA and RE commented on drafts of the manuscript.

COMPETING INTERESTS

AS is joint Editor-in-Chief of npjPCRM, but was not involved in the editorial review of, nor the decision to publish, this article. None of the other authors report any conflict of interest.

FUNDING

This study was supported by the Saudi Arabian Ministry of Higher Education Scholarship. AS is supported by The Commonwealth Fund, a private independent foundation based in New York City. The views presented here are those of the author and not necessarily those of The Commonwealth Fund, its directors, officers or staff.

REFERENCES

- Bereket A, Kiess W, Lustig RH, Muller HL, Goldstone AP, Weiss R *et al*. Hypothalamic obesity in children. *Obes Rev* 2012; **13**: 780–798.
- Santos JL, Ho-Urriola JA, Gonzalez A, Smalley SV, Domínguez-Vásquez P, Cataldo R *et al*. Association between eating behavior scores and obesity in Chilean children. *Nutr J* 2011; **10**: 108.
- Scopinaro N, Adami GF, Papadia FS, Camerini G, Carlini F, Briatore L *et al*. The effects of biliopancreatic diversion on type 2 diabetes mellitus in patients with mild obesity (BMI 30–35 kg/m²) and simple overweight (BMI 25–30 kg/m²): a prospective controlled study. *Obes Surg* 2011; **21**: 880–888.
- Badran M, Laher I. Obesity in Arabic-speaking countries. *J Obes* 2011; **2011**: 686430.
- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA* 2012; **307**: 483–490.

- 6 Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. *Lancet* 1998; **351**: 1225–1232.
- 7 Anandan C, Nurmatov U, van Schayck OC, Sheikh A. Is the prevalence of asthma declining? Systematic review of epidemiological studies. *Allergy* 2010; **65**: 152–167.
- 8 Anderson HR, Gupta R, Strachan DP, Limb ES. 50 years of asthma: UK trends from 1955 to 2004. *Thorax* 2007; **62**: 85–90.
- 9 Bustos P, Amigo H, Oyarzun M, Rona RJ. Is there a causal relation between obesity and asthma? Evidence from Chile. *Int J Obes* 2005; **29**: 804–809.
- 10 Shaheen SO. Obesity and asthma: cause for concern? *Clin Exp Allergy* 1999; **29**: 291–293.
- 11 Tantisira KG, Weiss ST. Complex interactions in complex traits: obesity and asthma. *Thorax* 2001; **56**(Suppl 2): ii64–ii73.
- 12 Camargo CA Jr, Weiss ST, Zhang S, Willett WC, Speizer FE. Prospective study of body mass index, weight change, and risk of adult-onset asthma in women. *Arch Intern Med* 1999; **159**: 2582–2588.
- 13 Nystad W, Meyer HE, Nafstad P, Tverdal A, Engeland A. Body mass index in relation to adult asthma among 135,000 Norwegian men and women. *Am J Epidemiol* 2004; **160**: 969–976.
- 14 Cassol VE, Rizzato TM, Teche SP, Basso DF, Centenaro DF, Maldonado M et al. Obesity and its relationship with asthma prevalence and severity in adolescents from southern Brazil. *J Asthma* 2006; **43**: 57–60.
- 15 Noal RB, Menezes AM, Macedo SE, Dumith SC. Childhood body mass index and risk of asthma in adolescence: a systematic review. *Obes Rev* 2010; **12**: 93–104.
- 16 Mamun AA, Lawlor DA, Alati R, O'Callaghan MJ, Williams GM, Najman JM. Increasing body mass index from age 5 to 14 years predicts asthma among adolescents: evidence from a birth cohort study. *Int J Obes* 2007; **31**: 578–583.
- 17 Vangeepuram N, Teitelbaum SL, Galvez MP, Brenner B, Doucette J, Wolff MS. Measures of obesity associated with asthma diagnosis in ethnic minority children. *J Obes* 2011; **2011**: 517417.
- 18 Kajbaf TZ, Asar S, Alipoor MR. Relationship between obesity and asthma symptoms among children in Ahvaz, Iran: a cross sectional study. *Ital J Pediatr* 2011; **37**: 1.
- 19 Mishra V. Effect of obesity on asthma among adult Indian women. *Int J Obes Relat Metab Disord* 2004; **28**: 1048–1058.
- 20 Camargo CAJ, Wentowski C, Field A, Gillman M, Frazier A, Colditz G. Prospective cohort study of body mass index and risk of asthma in children. *Ann Epidemiol* 2003; **13**: 565.
- 21 Matricardi PM, Gruber C, Wahn U, Lau S. The asthma-obesity link in childhood: open questions, complex evidence, a few answers only. *Clin Exp Allergy* 2007; **37**: 476–484.
- 22 Hill AB. The environment and disease: association or causation? *Proc R Soc Med* 1965; **58**: 295–300.
- 23 Shore SA. Obesity and asthma: lessons from animal models. *J Appl Physiol* 2007; **102**: 516–528.
- 24 Shore SA. Obesity and asthma: possible mechanisms. *J Allergy Clin Immunol* 2008; **121**: 1087–1093, quiz 94–5.
- 25 Hallstrand TS, Fischer ME, Wurfel MM, Afari N, Buchwald D, Goldberg J. Genetic pleiotropy between asthma and obesity in a community-based sample of twins. *J Allergy Clin Immunol* 2005; **116**: 1235–1241.
- 26 Weiss ST, Shore S. Obesity and asthma: directions for research. *Am J Respir Crit Care Med* 2004; **169**: 963–968.
- 27 Weiss ST. Obesity: insight into the origins of asthma. *Nat Immunol* 2005; **6**: 537–539.
- 28 Sood A. Sex differences: implications for the obesity-asthma association. *Exerc Sport Sci Rev* 2011; **39**: 48–56.
- 29 Deesomchok A, Fisher T, Webb KA, Ora J, Lam YM, Loughheed MD et al. Effects of obesity on perceptual and mechanical responses to bronchoconstriction in asthma. *Am J Respir Crit Care Med* 2010; **181**: 125–133.
- 30 Nahhas M, Bhopal R, Anandan C, Elton R, Sheikh A. Prevalence of allergic disorders among primary school-aged children in Madinah, Saudi Arabia: two-stage cross-sectional survey. *PLoS ONE* 2012; **7**: e36848.
- 31 Trasande L, Cronk C, Durkin M, Weiss M, Schoeller DA, Gall EA et al. Environment and obesity in the National Children's Study. *Environ Health Perspect* 2009; **117**: 159–166.
- 32 Baker KM, Goetzel RZ, Pei X, Weiss AJ, Bowen J, Tabrizi MJ et al. Using a return-on-investment estimation model to evaluate outcomes from an obesity management worksite health promotion program. *J Occup Environ Med* 2008; **50**: 981–990.
- 33 Sole D, Vanna AT, Yamada E, Rizzo MC, Naspitz CK. International Study of Asthma and Allergies in Childhood (ISAAC) written questionnaire: validation of the asthma component among Brazilian children. *J Investig Allergol Clin Immunol* 1998; **8**: 376–382.
- 34 Mohammad Y, Tabbah K, Mohammad S, Yassine F, Clayton T, Hassan M. International study of asthma and allergies in childhood: phase 3 in the Syrian Arab Republic. *East Mediterr Health J* 2010; **16**: 710–716.
- 35 Bener A, Abdulrazzaq YM, Debusse P, Al-Mutawwa J. Prevalence of asthma among Emirates school children. *Eur J Epidemiol* 1994; **10**: 271–278.
- 36 Janahi IA, Bener A, Bush A. Prevalence of asthma among Qatari schoolchildren: International Study of Asthma and Allergies in Childhood, Qatar. *Pediatr Pulmonol* 2006; **41**: 80–86.
- 37 Stumvoll M, Tschritter O, Fritsche A, Staiger H, Renn W, Weisser M et al. Association of the T-G polymorphism in adiponectin (exon 2) with obesity and insulin sensitivity: interaction with family history of type 2 diabetes. *Diabetes* 2002; **51**: 37–41.
- 38 Tanner JM, Whitehouse RH, Takaishi M. Standards from birth to maturity for height, weight, height velocity, and weight velocity: British children, 1965. II. *Arch Dis Child* 1966; **41**: 613–635.
- 39 Weiss M. Diabetes–renal failure–obesity–osteoporosis. How the thyroid gland controls metabolism. *MMW Fortschritte der Medizin* 2007; **149**: 10–12.
- 40 Durnin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr* 1974; **32**: 77–97.
- 41 Reilly JJ, Wilson J, Durnin JV. Determination of body composition from skinfold thickness: a validation study. *Arch Dis Child* 1995; **73**: 305–310.
- 42 Hatipoglu N, Ozturk A, Mazicioglu MM, Kurtoglu S, Seyhan S, Lokoglu F. Waist circumference percentiles for 7- to 17-year-old Turkish children and adolescents. *Eur J Pediatr* 2008; **167**: 383–389.
- 43 Miller MR, Crapo R, Hankinson J, Brusasco V, Burgos F, Casaburi R et al. General considerations for lung function testing. *Eur Respir J* 2005; **26**: 153–161.
- 44 Clausen JL, Coates AL, Quanjer PH. Measurement of lung volumes in humans: review and recommendations from an ATS/ERS workshop. *Eur Respir J* 1997; **10**: 1205–1206.
- 45 Beydon N, Davis SD, Lombardi E, Allen JL, Arets HG, Aurora P et al. An official American Thoracic Society/European Respiratory Society statement: pulmonary function testing in preschool children. *Am J Respir Crit Care Med* 2007; **175**: 1304–1345.
- 46 Arshad SH, Tariq SM, Matthews S, Hakim E. Sensitization to common allergens and its association with allergic disorders at age 4 years: a whole population birth cohort study. *Pediatrics* 2001; **108**: E33.
- 47 Bryant T. *Confidence Interval Analysis (CIA)*, version 2 edn. BMJ Publishing Group: London, UK.
- 48 Castro-Rodriguez JA, Holberg CJ, Morgan WJ, Wright AL, Martinez FD. Increased incidence of asthmalike symptoms in girls who become overweight or obese during the school years. *Am J Respir Crit Care Med* 2001; **163**: 1344–1349.
- 49 Hancox RJ, Milne BJ, Poulton R, Taylor DR, Greene JM, McLachlan CR et al. Sex differences in the relation between body mass index and asthma and atopy in a birth cohort. *Am J Respir Crit Care Med* 2005; **171**: 440–445.
- 50 Magnusson JO, Kull I, Mai XM, Wickman M, Bergström A. Early childhood overweight and asthma and allergic sensitization at 8 years of age. *Pediatrics* 2012; **129**: 70–76.
- 51 Kannan S, Acosta LM, Acevedo-Garcia D, Divjan A, Bracero LA, Perzanowski MS et al. Sociocultural characteristics, obesity and inflammatory biomarkers in Puerto Rican toddlers born in New York City. *Pediatr Allergy Immunol* 2013; **24**: 487–492.
- 52 Hersoug LG, Linneberg A. The link between the epidemics of obesity and allergic diseases: does obesity induce decreased immune tolerance? *Allergy* 2007; **62**: 1205–1213.
- 53 Guidotti TL. Applying the precautionary principle. *Arch Environ Occup Health* 2012; **67**: 63–64.
- 54 Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr* 2007; **150**: 12–7 e2.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0>